


MULTI-ENGINEERING OF MICROBIAL CYTOCHROME P450 ENZYMES

Shengying Li, Shandong University, China; Shandong Provincial Key Laboratory of Synthetic Biology, Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, China
lishengying@sdu.edu.cn

Key Words: Cytochrome P450 enzyme, Enzyme engineering, Substrate engineering, Redox partner engineering, Electron donor engineering

Selective oxidation of unactivated C–H bonds remains a central challenge in synthetic chemistry. Cytochrome P450 enzymes, a superfamily of ubiquitous hemoproteins, represent the nature's primary solutions to overcome this challenge. As promising biocatalysts for practical applications in pharmaceutical, biotechnological and chemical industries, P450 enzymes have attracted a wealth of attention due to their great versatility in catalyzing diverse oxidative reactions (e.g., the sp^3 C–H hydroxylation and the sp^2 C=C epoxidation) on structurally complex and heavily functionalized substrates in regio- and/or stereoselective manners. However, wild type P450 enzymes usually show suboptimal activity, low stability, and narrow substrate spectra, which have significantly limited their broader applications. A typical P450 reaction system includes a P450 enzyme as the central catalyst, a substrate to be oxidized, redox partner proteins for electron transfer, NAD(P)H as the electron donor, and O_2 as the oxidant. In the past five years, we have made significant progresses on enzyme engineering, substrate engineering, redox partner engineering, and electron donor engineering for a number of microbial P450 enzymes. These multi-engineering efforts have generated useful engineered P450 catalytic systems for bio-production of pharmaceuticals, chemical intermediates, and biofuel molecules.



Reference:

1. Zhang, W., Du, L., Li, F., Zhang, X., Qu, Z., Han, L., Li, Z., Sun, J., Qi, F., Yao, Q., Sun, Y., Geng, C. and Li, S. Mechanistic Insights into Interactions between Bacterial Class I P450 Enzymes and Redox Partners. *ACS Catal.* 2018, 8, 9992-10003.
2. Du, L., Dong, S., Zhang, X., Jiang, C., Chen, J., Yao, L., Wang, X., Wan, X., Liu, X., Wang, X., Huang, S., Cui, Q., Feng, Y., Liu, S., and Li, S. Selective oxidation of aliphatic C–H bonds in alkylphenols by a chemomimetic biocatalytic system. *Proc. Natl. Acad. Sci. U.S.A.* 2017, 114, E5129-E5137.
3. Ma, L., Du, L., Chen, H., Sun, Y., Huang, S., Zheng, X., Kim, E.-S., and Li, S. Reconstitution of the in vitro activity of the cyclosporine-specific P450 hydroxylase from *Sebekia benihana* and development of a heterologous whole cell biotransformation system. *Appl. Environ. Microbiol.* 2015, 81, 6268-6275.
4. Zhang, W., Liu, Y., Yan, J., Cao, S., Bai, F., Yang, Y., Huang, S., Yao, L., Anzai, Y., Kato, F., Podust, L. M., Sherman, D. H., and Li, S. New reactions and products resulting from alternative interactions between the P450 enzyme and redox partners. *J. Am. Chem. Soc.* 2014, 136, 3640-3646.
5. Liu, Y., Wang, C., Yan, J., Zhang, W., Guan, W., Lu, X., and Li, S. Hydrogen peroxide-independent production of α -alkenes by OleT_{JE} P450 fatty acid decarboxylase. *Biotechnol. Biofuels* 2014, 7, 28.